

Chiral Symmetry Restoration and Eigenvalue Density of Dirac Operator at Finite Temperature

Sinya AOKI

University of Tsukuba

with N. Fukaya and Y. Taniguchi for JLQCD Collaboration

“New Horizon for Lattice Gauge Theory Computations with Chiral Fermions ”,
2012.05.14-16, BNL, NY, USA

1. Introduction

Flavor-Chiral Symmetries of QCD at low T

$$\begin{array}{ccccc}
 U(N_f)_L \otimes U(N_f)_R & \xrightarrow{\text{chiral anomaly (explicit breaking)}} & U(1)_B \otimes S(N_f)_L \otimes SU(N_f)_R & \xrightarrow{\text{spontaneous breaking of chiral symmetry}} & U(1)_B \otimes S(N_f)_V \\
 \text{flavor} & & & &
 \end{array}$$

QCD at high T

deconfinement (QGP)

restoration of chiral symmetry

phase transition

$$\begin{array}{ccccc}
 \text{low T} & U(1)_B \otimes S(N_f)_V & \xrightarrow{\text{phase transition}} & \text{high T} & U(1)_B \otimes S(N_f)_L \otimes SU(N_f)_R
 \end{array}$$

Questions in this study

1. Constraints to eigenvalue density:

$$\rho(\lambda) = \lim_{V \rightarrow \infty} \frac{1}{V} \sum_n \delta(\lambda - \lambda_n)$$

2. Constraint to singlet susceptibility:










$$\chi_{\text{singlet}} = \int d^4x [\pi(x)\pi(0) - \eta(x)\eta(0)]$$

Previous studies

$$\rho(\lambda) = \lim_{V \rightarrow \infty} \frac{1}{V} \sum_n \delta(\lambda - \lambda_n)$$

Lin (HotQCD11), DW


Cossu *et al.* (JLQCD11), Overlap




Ohno *et al.* (11), HISQ

Is a gap open at $\lambda = 0$?


Chandrasekharan-Christ(95), KS




Bernald, *et al.* (96), KS



Chandrasekharan *et al.*, (98), KS



Hegde (HotQCD11), DW



$\chi_{\text{singlet}} = 0 ?$

This talk

give constraints to eigenvalue densities of **2-flavor overlap fermions**, if chiral symmetry in QCD is restored at finite temperature.

discuss a behavior of singlet susceptibility using the constraints.

Content

1. Introduction
2. Overlap fermions
3. Constraints to eigenvalue densities
4. Discussions: singlet susceptibility

2. Overlap fermions


Action $S = \bar{\psi}[D - mF(D)]\psi, \quad F(D) = 1 - \frac{Ra}{2}D$

Ginsparg-Wilson relation $D\gamma_5 + \gamma_5 D = aDR\gamma_5 D$


Eigenvalue spectrum

$$\lambda_n^A + \bar{\lambda}_n^A = aR\bar{\lambda}_n^A \lambda_n^A$$

$$D(A)\phi_n^A = \lambda_n^A \phi_n^A$$



$$D(A)\gamma_5 \phi_n^A = \bar{\lambda}_n^A \gamma_5 \phi_n^A$$



Propagator

bulk modes(non-chiral)

zero modes(chiral)

doublers(chiral)

$$S(x, y) = \sum_n \left[\frac{\phi_n(x)\phi_n^\dagger(y)}{f_m \lambda_n - m} + \frac{\gamma_5 \phi_n(x)\phi_n^\dagger(y)\gamma_5}{f_m \bar{\lambda}_n - m} \right] - \sum_{k=1}^{N_{R+L}} \frac{1}{m} \phi_k(x)\phi_k^\dagger(y) + \sum_{K=1}^{N_D} \frac{Ra}{2} \phi_K(x)\phi_K^\dagger(y)$$

$$f_m = 1 + \frac{Rma}{2}$$

Measure $P_m(A) = e^{-S_{YM}(A)} (-m)^{N_f N_{R+L}^A} \left(\frac{2}{Ra} \right)^{N_f N_D^A} \prod_{\Im \lambda_n^A > 0} (Z_m^2 \bar{\lambda}_n^A \lambda_n^A + m^2)$


$$Z_m^2 = 1 - (ma)^2 \frac{R^2}{4}$$

positive definite and even function of $m \neq 0$ for even N_f

Ward-Takahashi identities under “chiral” rotation

$$\begin{aligned}\theta^a(x)\delta_x^a\psi(x) &= i\theta^a(x)T^a\gamma_5(1-RaD)\psi(x), \\ \theta^a(x)\delta_x^a\bar{\psi}(x) &= i\bar{\psi}(x)\theta^a(x)T^a\gamma_5,\end{aligned}$$

$$\langle (J_x^a - \delta_x^a S)O + \delta_x^a \mathcal{O} \rangle = 0$$

 $\int d^4x \langle \{J_x^a + 2mP^a(x)\}\mathcal{O} + \delta_x^a \mathcal{O} \rangle = 0.$ integrated WTI

“measure” term (anomaly)

$$J_x^a = -2i\text{tr} T^a\gamma_5 \left(1 - \frac{R}{2}aD\right)(x,x) = -\delta^{a0}2iN_f\text{tr} \gamma_5 \left(1 - \frac{R}{2}aD\right)(x,x)$$

Operators

$$\begin{aligned}S^a(x) &= \bar{\psi}(x)T^aF(D)\psi(x), && \text{scalar} \\ P^a(x) &= \bar{\psi}(x)T^ai\gamma_5F(D)\psi(x), && \text{pseudo-scalar}\end{aligned} \qquad S^a = \int d^4x S^a(x), \quad P^a = \int d^4x P^a(x)$$

chiral rotation at N_f=2

$$\begin{aligned}\delta^b S^a &= 2\delta^{ab}P^a, & \delta^b P^a &= -2\delta^{ab}S^a \\ \delta^0 S^a &= \delta^a S^0 = 2P^a, & \delta^0 P^a &= \delta^a P^0 = -2S^a\end{aligned}$$

$$\mathcal{O}_{n_1,n_2,n_3,n_4} = (P^a)^{n_1}(S^a)^{n_2}(P^0)^{n_3}(S^0)^{n_4} \qquad N = \sum_i n_i, \quad n_1 + n_2 = \text{odd}, \quad n_1 + n_3 = \text{odd}$$

$\lim_{m \rightarrow 0} \langle \delta^a \mathcal{O}_{n_1,n_2,n_3,n_4} \rangle_m = 0$

if the chiral symmetry is restored.

$$\frac{\delta^a}{2}\mathcal{O}_{n_1,n_2,n_3,n_4} = -n_1\mathcal{O}_{n_1-1,n_2,n_3,n_4+1} + n_2\mathcal{O}_{n_1,n_2-1,n_3+1,n_4} - n_3\mathcal{O}_{n_1,n_2+1,n_3-1,n_4} + n_4\mathcal{O}_{n_1+1,n_2,n_3,n_4-1}$$

3. Constraints to eigenvalue densities

$$\boxed{N=1} \quad \mathcal{O}_{1,0,0,0} = P^a, \quad \delta^a \mathcal{O}_{1,0,0,0} = -2S^0$$

$$\frac{1}{V} \langle S^0 \rangle_m = \frac{N_f}{mV} \langle \underline{N_{R+L}^A} \rangle_m + N_f \langle I_1 \rangle_m \qquad \langle \mathcal{O}^A \rangle_m = \frac{1}{Z} \int \mathcal{D}A P_m(A) \mathcal{O}^A$$

$$\rightarrow 0 \text{ as } V \rightarrow \infty$$

$$I_1 \rightarrow \frac{1}{Z_m} \int_0^{\Lambda_R} d\lambda \underline{\rho^A(\lambda)} \frac{2m_R}{\lambda^2 + m_R^2} g_0(\lambda)$$

$$m_R = \frac{m}{Z_m}, \quad g(\lambda) = 1 - \frac{\lambda^2}{\Lambda_R^2}$$

$$\Lambda_R = \frac{2}{Ra}: \text{ cut-off}$$

$$\rho^A(\lambda) \equiv \lim_{V \rightarrow \infty} \frac{1}{V} \sum_n \delta \left(\lambda - \sqrt{\bar{\lambda}_n^A \lambda_n^A} \right) = \sum_{k=0}^{\infty} \rho_k^A \frac{|\lambda|^k}{k!} = \rho_0^A + \rho_1^A |\lambda| + \dots,$$

$$\Rightarrow I_1 = \pi \rho_0^A + O(m)$$

$$\lim_{m \rightarrow 0} \lim_{V \rightarrow \infty} \frac{1}{V} \langle S^0 \rangle_m = 0 \Rightarrow \lim_{m \rightarrow 0} \langle \rho_0^A \rangle_m = 0$$

$$\Rightarrow \langle \rho_0^A \rangle_m = O(m^2) \quad \text{1st constraint}$$

$$\boxed{N=2}$$

$$\chi^{\sigma-\pi} = \frac{1}{V^2} \langle S_0^2 - P_a^2 \rangle_m, \quad \chi^{\eta-\delta} = \frac{1}{V} \langle P_0^2 - S_a^2 \rangle_m$$

$$\chi^{\sigma-\pi} = \left\langle N_f^2 \left(\frac{1}{mV} N_{R+L} + I_1 \right)^2 \right\rangle_m + O\left(\frac{1}{V}\right)$$

$$\rightarrow \lim_{m \rightarrow 0} \lim_{V \rightarrow \infty} \frac{1}{m^2 V^2} \underbrace{\langle (N_{R+L}^A)^2 \rangle_m}_{O(m^{N_f V})} = \lim_{m \rightarrow 0} \lim_{V \rightarrow \infty} \frac{1}{mV} \underbrace{\langle N_{R+L}^A \rho_0^A \rangle_m}_{O(m^{N_f V})} = \lim_{m \rightarrow 0} \underbrace{\langle (\rho_0^A)^2 \rangle_m}_{O(m^2)} = 0$$

These conditions are automatically satisfied.

$$\chi^{\eta-\delta} = N_f \left\langle \frac{1}{m^2 V} \{2N_{R+L} - N_f Q(A)^2\} + \frac{1}{Z_m} \left(\frac{I_1}{m_R} + I_2 \right) \right\rangle_m \quad Q(A) = N_R^A - N_L^A$$

$$\frac{I_1}{m_R} + I_2 = \rho_0^A \left(\frac{\pi_m}{m} + \frac{2}{\Lambda_R} \right) + 2\rho_1^A + O(m), \quad I_2 = \frac{2}{Z_m} \int_0^{\Lambda_R} d\lambda \rho^A(\lambda) \frac{m_R^2 - \lambda^2 g_0(\lambda^2) g_m}{(\lambda^2 + m_R^2)^2}, \quad g_m = \frac{1}{Z_m^2} \left(1 + \frac{m^2}{2\Lambda_R^2} \right)$$

$$\lim_{m \rightarrow 0} \chi^{\eta-\delta} = 0 \rightarrow \lim_{m \rightarrow 0} \underbrace{\frac{N_f^2 \langle Q(A)^2 \rangle_m}{m^2 V}}_{O(m^{N_f \sqrt{V}-2})} = 2 \lim_{m \rightarrow 0} \langle \rho_1^A \rangle_m$$

$$\rightarrow \langle \rho_1^A \rangle_m = O(m^2) \quad \text{2nd constraint}$$


N=3

$$\begin{aligned} \langle \mathcal{O}_{2001} \rangle_m &\rightarrow 0, & \langle -\mathcal{O}_{0201} + 2\mathcal{O}_{1110} \rangle_m &\rightarrow 0, & \langle \mathcal{O}_{0021} + 2\mathcal{O}_{1110} \rangle_m &= 0 \\ \langle -\mathcal{O}_{0003} + 2\mathcal{O}_{2001} \rangle_m &\rightarrow 0, & \langle \mathcal{O}_{0021} - \mathcal{O}_{0201} + \mathcal{O}_{1110} \rangle_m &\rightarrow 0, \end{aligned}$$

$$\lim_{m \rightarrow 0} \lim_{V \rightarrow \infty} \frac{\chi_3}{V^4} = \pi N_f \lim_{m \rightarrow 0} \left[\frac{\langle \rho_0^A \rangle_m}{m^2} + \frac{2\bar{\rho}_1}{\underline{m\pi}} + \frac{\langle \rho_2^A \rangle_m}{2} \right], = 0$$

+ positivity

$O(m)$

 $\langle \rho_0^A \rangle_m = O(m^4), \quad \langle \rho_2^A \rangle_m = O(m^2)$ 3rd constraints

N=4

No additional constraints

Final results

$$\lim_{m \rightarrow 0} \langle \rho^A(\lambda) \rangle_m = \lim_{m \rightarrow 0} \langle \rho_3^A \rangle_m \frac{|\lambda|^3}{3!} + O(\lambda^4)$$

4. Discussion: Singlet susceptibility

Singlet susceptibility

$$\chi^{\pi-\eta} = \lim_{V \rightarrow \infty} \frac{N_f^2}{m^2 V} \langle Q(A)^2 \rangle_m = (m^{N_f \sqrt{V}-2})$$

➔ $\lim_{m \rightarrow 0} \chi^{\pi-\eta} = 0$

singlet susceptibility becomes zero if the chiral symmetry is recovered at high T.

This, however, does not mean $U(1)_A$ symmetry is recovered at high T.

$\lim_{m \rightarrow 0} \chi^{\pi-\eta} = 0$ is necessary but NOT sufficient for the recovery of $U(1)_A$.

Future

new constraints at $N > 4$?

eigenvalues of Dirac operator
have a gap near zero ?

$$\rho(\lambda) = 0 \text{ at } |\lambda| \leq \lambda_c$$

